

2/13




Figure 2: Synthesizing the hydrophobic bicatenary hydrocarbon and perfluorocarbon moieties

3/13





Figure 3: Synthesizing the nonionic polar heads






Figure 5: Synthesizing the monocatenary amphiphilic nitrones **A-B**




Figure 6: Synthesizing the bicatenary amphiphilic nitrones C-D

7/13




Figure 7: EPR spectra of the carboxylate (a), hydroxyl (b) and methyl (c) adducts which are respectively generated by the Fenton reaction (b) in the presence of formate (a) and of DMSO (c) and of the compound **A₁**




Figure 8: Caspase III activity of neuronal cells which have been poisoned with H₂O₂ and treated with commercial nitrones and the type **A2** nitrosonium






Figure 9: Measuring the state of apoptosis by means of an ELISA assay of the fragmentation of the DNA following lysis of the cells

10/13



Figures 10a and 10b: Culture of NARP cells in the presence of 100 (a) and 200 μ M (b) amphiphilic nitrones after 48 h of incubation


11/13




Figures 10c and 10d: Culture of NARP cells in the presence of 100 (d) and 200 μ M (c) amphiphilic nitrones after 72 h of incubation

12/13

e



f



Figures 10e and 10f: Culture of NARP cells in the presence of 50 μ M amphiphilic nitrones after 48 h (e) and 72 h (f) of incubation




Figure 11: Variation in the hydrophobicity of the lactobionic acid-derived nitrones **A1-A5**.